

Title	THE VERTICES OF THE COMPONENTS OF THE PERMUTATION MODULE INDUCED FROM PARABOLIC GROUPS
Author(s)	Pforte, Lars
Citation	Osaka Journal of Mathematics. 55(4) p.769-p.775
Issue Date	2018-10
oaire:version	VoR
URL	https://doi.org/10.18910/70821
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

THE VERTICES OF THE COMPONENTS OF THE PERMUTATION MODULE INDUCED FROM PARABOLIC GROUPS

LARS PFORTE

(Received October 25, 2016, revised June 16, 2017)

Abstract

We consider the permutation module $k_P \uparrow^{\mathrm{GL}_n(p^f)}$, where P is a parabolic group in the general linear group $\mathrm{GL}_n(p^f)$ and k is an algebraically closed field of prime characteristic p . The vertices of the components of these modules have been calculated in [9] by Tinberg, who studied these modules for all groups with split BN-pairs in characteristic p . In this paper we show that the idea of suitability is strong enough to find all p -groups that are vertex of some component of $k_P \uparrow^{\mathrm{GL}_n(p^f)}$. Furthermore using a result of Burry and Carlson we show that all components have a different vertex.

1. Introduction

Let G be a finite group with subgroup H . We are interested in the p -groups of G that are vertices of components of the permutation-module $k_H \uparrow^G$, where k is an algebraically closed field of characteristic p . In [7] the author introduces the idea of H -suitability and in an example shows how suitability could be applied to detect all p -groups that are vertices of some component of $k_H \uparrow^G$. Furthermore in that example suitability is at its most restrictive as every H -suitable group turns out to be the vertex of a component of $k_H \uparrow^G$. In the present paper we present another example where this concept of suitability unfolds its full potential by giving us only p -groups that are vertices of some component of the given permutation module $k_H \uparrow^G$.

Next we describe the notion of H -suitability. A p -subgroup V of G is called H -suitable (with respect to G and p), if for every $S \in \mathrm{Syl}_p(G)$ with $V \leq S$ there exists some $g \in G$ so that $V = S \cap H^g$. Here H^g denotes the conjugate $g^{-1}Hg$ of H . In [7] we show that H -suitability is a necessary condition for a p -group to be the vertex of a component of $k_H \uparrow^G$ and present an example in which it is also sufficient. However in general an H -suitable group is not necessarily the vertex of some component of $k_H \uparrow^G$. For instance with respect to the symmetric group S_5 and $p = 2$ the trivial group is D -suitable, for the dihedral group $D := \langle (1234), (13) \rangle \leq S_5$, but $k_D \uparrow^{S_5}$ has no projective summand. Nevertheless one can use the concept of suitability to produce a list of potential candidates for a vertex by finding all H -suitable groups. If this list is short enough one may then deal with each candidate individually.

In [7] we see that if V is H -suitable then so is every G -conjugate of V . Hence given a Sylow- p -group S of G the set $\mathcal{A} := \{S \cap H^g : g \in G\}$ contains a representative of each

conjugacy class of H -suitable groups. The following statement presents a method to shorten the list given by \mathcal{A} by finding groups that are not H -suitable. A more general version of the statement with proof can be found in [7, Theorem 2.6.].

Lemma 1.1. *Let $S \in \text{Syl}_p(G)$ and $K \leq G$. Also let $T \subseteq G$ such that*

- (a) *For every $g \in G$ there is $\alpha \in T$ so that $S \cap H^g =_K S \cap H^{\alpha^{-1}}$.*
- (b) *For all $\alpha \in T$ we have $S^\alpha \cap H \leq S$.*
- (c) *If $S^\alpha \cap H =_K S^{\alpha'} \cap H$, for $\alpha, \alpha' \in T$, then $\alpha = \alpha'$.*

Furthermore let $V \leq S$ be H -suitable. Then there is some $\alpha \in T$ such that $V =_K S^\alpha \cap H$.

Observe that Lemma 1.1 implies that every H -suitable group V that is contained in S is a K -conjugate of both $S \cap H^{\alpha^{-1}}$ and $S^\beta \cap H$, for some $\alpha, \beta \in T$. In particular any group in the set $\{S \cap H^{\alpha^{-1}} : \alpha \in T\}$ which has no K -conjugate in the set $\{S^\alpha \cap H : \alpha \in T\}$ fails to be H -suitable.

Let us now turn to our example for which we need the following notation. For the remainder of this paper let $n, f \geq 1$ and set $q = p^f$, for some prime number p . Furthermore let $G := \text{GL}_n(q)$ be the general linear group, that is, the group of invertible $n \times n$ -matrices over the finite field \mathbb{F}_q with q elements. Finally let k be an algebraically closed field of characteristic p .

For any integer $j \geq 1$ we define $\text{GL}_j := \text{GL}_j(q)$. By B_j we denote the group of upper-triangular matrices in GL_j . Those elements in B_j with 1's on the main diagonal form the group U_j . Recall that U_j is a Sylow- p -group of GL_j and B_j is the normalizer of U_j in GL_j .

Furthermore \mathcal{W}_j denotes the group of permutation matrices in GL_j . There is a bijection between the symmetric group $\text{Sym}(j)$ and \mathcal{W}_j , in the sense that the permutation $\omega \in \text{Sym}(j)$ corresponds to the permutation matrix $(\delta_{r, \omega(s)})_{r,s}$, where δ denotes the Kronecker-symbol.

Let $\lambda = (\lambda_1, \dots, \lambda_r)$ be a *composition* of n , that is, $\lambda_1, \dots, \lambda_r$ are positive integers such that $\lambda_1 + \dots + \lambda_r = n$. Then for $X_t \in \text{GL}_{\lambda_t}$, let $D_n(X_1, X_2, \dots, X_r)$ be the matrix in G , which has X_1, \dots, X_r along the main diagonal and zeros otherwise. Similarly $D_n(X_1 \bullet, X_2 \bullet, \dots, X_r)$ denotes any matrix in G which has X_1, \dots, X_r along the main diagonal, arbitrary elements in \mathbb{F}_q above and zeros below that diagonal. Finally we define

$$P_\lambda = \{D_n(X_1 \bullet, X_2 \bullet, \dots, X_r) : X_t \in \text{GL}_{\lambda_t}\}.$$

Then P_λ is a *parabolic subgroup* of G . For instance if $\lambda = (1^n)$ then $P_\lambda = B_n$.

The permutation module $k_{P_\lambda} \uparrow^G$, where λ is a composition of n , has been studied before and in more generality. Tinberg [9] studied these modules for all groups with split BN-pairs in characteristic p . Further work was done by Canbanes [3] and Sin [8]. In particular the vertices of all components are known as well as their Green correspondents. In this paper we revisit $k_{P_\lambda} \uparrow^G$ and show that P_λ -suitability is a strong enough tool to find all p -groups that are vertices of components of $k_{P_\lambda} \uparrow^G$. In section 2 we find all P_λ -suitable groups up to G -conjugation and in section 3 we show that all P_λ -suitable groups are vertex of exactly one component of $k_{P_\lambda} \uparrow^G$.

Next let $1 \leq k, l \leq n$ such that $k \neq l$. By $F_{k,l}$ we denote the subgroup of G that consists of exactly those matrices that have ones on the main diagonal and zeros everywhere else except in the (k, l) -entry, which is arbitrary in \mathbb{F}_q . One checks easily that for $\omega \in \mathcal{W}_n$ and $k \neq l$ we

have $F_{k,l} = (F_{\omega(k),\omega(l)})^\omega$. Finally by [5] a group $V \leq U_n$ is called a *pattern group* if $X_{k,l} \neq 0$, for some $X \in V$ and $1 \leq k < l \leq n$, implies that $F_{k,l} \leq V$. Note that a pattern group which contains both $F_{k,l}$ and $F_{l,m}$, for $k < l < m$, also contains $F_{k,m}$.

Lemma 1.2. *Let $V, W \leq U_n$ be pattern groups that are B_n -conjugate. Then $V = W$.*

Proof. It is enough to show that $F_{k,l} \leq V$ implies $F_{k,l} \leq W$. So let $F_{k,l} \leq V$, for some $k < l$, and let $g \in F_{k,l}$ be non-trivial. By assumption there is $h \in B_n$ so that $g' := g^h \in W$. As $g'_{k,l} \neq 0$ it now follows that $F_{k,l} \leq W$. \square

2. P_λ -suitable groups

In this section let $\lambda = (\lambda_1, \dots, \lambda_r)$ be a composition of n . Also set $s_i := \sum_{k=1}^{i-1} \lambda_k$, for $i = 1, \dots, r$, and $s_{r+1} := n$.

2.1. Good Permutations. We call $\omega \in \mathcal{W}_n$ *good* with respect to λ , if $\omega(s_i + 1) < \dots < \omega(s_i + \lambda_i)$, for all $i = 1, \dots, r$.

Lemma 2.1. *Every (B_n, P_λ) -double coset in G contains a permutation matrix in \mathcal{W}_n that is good with respect to λ .*

Proof. Let $g \in G$. Then by the Bruhat decomposition, (see [1]), and since $B_n \leq P_\lambda$, there exists $\mu \in \mathcal{W}_n$ such that $B_n \cdot g \cdot P_\lambda = B_n \cdot \mu \cdot P_\lambda$. Next let

$$\mu(\{s_i + 1, \dots, s_i + \lambda_i\}) = \{k_{s_i+1}, \dots, k_{s_i+\lambda_i}\}, \text{ where } k_{s_i+1} < \dots < k_{s_i+\lambda_i},$$

for all $i = 1, \dots, r$. Set $\gamma(j) := \mu^{-1}(k_j)$, for all $j = 1, \dots, n$. Then $\gamma \in \mathcal{W}_n$. Also observe that γ acts on the sets $\{s_i + 1, \dots, s_i + \lambda_i\}$. Hence $\gamma \in P_\lambda$.

Next set $\omega := \mu \cdot \gamma \in \mathcal{W}_n$. Since $\omega(j) = \mu(\mu^{-1}(k_j)) = k_j$, for all $j = 1, \dots, n$, it follows that ω is good. As now $B_n \cdot g \cdot P_\lambda = B_n \cdot \omega \cdot P_\lambda$, the proof is complete. \square

For any $g \in G$ we define $V(g) := U_n^g \cap P_\lambda$. The following lemma is then an easy exercise.

Lemma 2.2. *Let $\omega \in \mathcal{W}_n$ be good with respect to λ . Then*

$$(1) \quad \begin{aligned} V(\omega) &= \{X = D_n(A_1 \bullet, A_2 \bullet, \dots, A_r) : A_i \in U_{\lambda_i}, \text{ and} \\ &\quad X_{k,l} = 0, \text{ for all } k, l = 1, \dots, n \text{ so that } \omega(k) > \omega(l)\}. \end{aligned}$$

In particular $V(\omega) \leq U_n$, and $V(\omega)$ is a pattern group.

Observe that Lemma 2.1 implies the following

Corollary 2.3. *Every P_λ -suitable group is G -conjugate to some $V(\omega)$, where $\omega \in \mathcal{W}_n$ is good with respect to λ .*

Lemma 2.4. *The set T of all $\omega \in \mathcal{W}_n$ that are good with respect to λ satisfies the properties (a)-(c) in Lemma 1.1, where $H = P_\lambda$, $S = U_n$ and $K = B_n$.*

Proof. Property (a) is a consequence of Lemma 2.1, property (b) follows since $V(\omega) \leq U_n$ and property (c) is a consequence of Lemma 1.2 and the structure of $V(\omega)$, as given in (1). \square

Proposition 2.5. *Let $V \leq U_n$ be a pattern group. If V is P_λ -suitable, then $V = V(\omega)$, where $\omega \in \mathcal{W}_n$ is good with respect to λ .*

Proof. By Lemma 2.4 and Lemma 1.1 we have $V =_{B_n} V(\omega)$, where $\omega \in \mathcal{W}_n$ is good with respect to λ . Now $V = V(\omega)$ follows from Lemma 1.2. \square

2.2. λ -permutations. Recall that I_j denotes the identity matrix in GL_j . By \overline{I}_j we mean the permutation matrix in GL_j that has all its ones on the anti-diagonal. Also let $\omega = D_n(K_1, \dots, K_u) \in \mathcal{W}_n$, where $K_i \in \{I_{n_i}, \overline{I}_{n_i}\}$, and $n_1 + \dots + n_u = n$. We call such ω a λ -permutation if for all $t = 1, \dots, r$ so that $\lambda_t \geq 2$ we have that ω acts on each of the following sets

$$\{1, \dots, s_t + 1\}, \{s_t + 2, \dots, s_{t+1} - 1\}, \{s_{t+1}, \dots, n\}.$$

Observe that every λ -permutation is good with respect to λ . In the following we show that if $\omega \in \mathcal{W}_n$ is a λ -permutation, then $V(\omega)$ is P_λ -suitable.

Lemma 2.6. *Let $\omega \in \mathcal{W}_n$ be a λ -permutation. Then*

$$V(\omega) = \{D_n(A_1 \bullet, A_2 \bullet, \dots, A_u) : \text{where } A_i \in U_{n_i}, \text{ if } K_i = I_{n_i}, \\ \text{and } A_i = I_{n_i}, \text{ if } K_i = \overline{I}_{n_i}\}.$$

Proof. Let S be the set on the right hand side. Then $S = U_n^\omega \cap U_n \leq P_\lambda$. Since $V(\omega) \leq U_n$ we get $S \leq U_n^\omega \cap P_\lambda = V(\omega) \leq U_n^\omega \cap U_n = S$. Thus $V(\omega) = S$. \square

Lemma 2.6 implies the following

Corollary 2.7. *Let $\omega \in \mathcal{W}_n$ be a λ -permutation. Then*

- (1) $B_n \subseteq N_G(V(\omega))$
- (2) *If $\mu \in \mathcal{W}_n$ such that $V(\omega)^\mu \leq U_n$, then $\mu = D_n(\mu_1, \dots, \mu_u)$ where $\mu_i = I_{n_i}$, if $K_i = I_{n_i}$, and $\mu_i \in \mathcal{W}_{n_i}$, if $K_i = \overline{I}_{n_i}$.*

Lemma 2.8. *Let $\omega \in \mathcal{W}_n$ be a λ -permutation. Also let N be the set of all matrices of the form $D_n(A_1 \bullet, A_2 \bullet, \dots, A_u)$ where $A_i \in B_{n_i}$, if $K_i = I_{n_i}$, and $A_i \in \text{GL}_{n_i}$, if $K_i = \overline{I}_{n_i}$. Moreover let $g \in G$ such that $V(\omega)^g \leq U_n$. Then $g \in N$. In particular $N_G(V(\omega)) = N$.*

Proof. By the Bruhat decomposition there are $A, B \in B_n$ and $\mu \in \mathcal{W}_n$ such that $g = A\mu B$. As $B_n \subseteq N_G(V(\omega))$ we have $V(\omega)^\mu \leq U_n$, and thus $\mu \in N$, by Corollary 2.7. Since $B_n \subseteq N$, we get $g \in N$. Now $N_G(V(\omega)) \subseteq N$ follows. As one checks easily that N normalizes $V(\omega)$, the proof is complete. \square

Corollary 2.9. *Let $\omega_1, \omega_2 \in \mathcal{W}_n$ be λ -permutations so that $V(\omega_1) =_G V(\omega_2)$. Then $\omega_1 = \omega_2$.*

Proof. As $V(\omega_1)^g = V(\omega_2) \leq U_n$, for some $g \in G$, we have $g \in N_G(V(\omega_1))$, by Lemma 2.8. Thus $V(\omega_1) = V(\omega_2)$. Now $\omega_1 = \omega_2$ follows from (1). \square

Proposition 2.10. *Let $\omega \in \mathcal{W}_n$ be a λ -permutation. Then $V(\omega)$ is P_λ -suitable.*

Proof. Since $V(\omega)^{\omega^{-1}} \leq U_n$, it follows $\omega^{-1} \in N_G(V(\omega))$, by Lemma 2.8. Thus $V(\omega) = U_n \cap P_\lambda^{\omega^{-1}}$. Note that $U_n \in \text{Syl}_p(N_G(V(\omega)))$. So the statement follows from [7, Lemma 2.2.(3)]. \square

2.3. The P_λ -suitable $V(\omega)$. In the following let $\omega \in \mathcal{W}_n$ be good with respect to λ so that $V(\omega)$ is P_λ -suitable. We aim to show that ω is a λ -permutation.

Lemma 2.11. *Let $F_{k,l} \leq V(\omega)$, that is, $\omega(k) < \omega(l)$, for some $k < l$. Then*

- (1) $F_{t,l} \leq V(\omega)$, that is, $\omega(t) < \omega(l)$, for all $t = 1, \dots, k$
- (2) $F_{k,t} \leq V(\omega)$, that is, $\omega(k) < \omega(t)$, for all $t = l, \dots, n$.

Proof. We prove part (1) by contradiction. Without loss of generality we may assume that $F_{k-1,l} \not\leq V(\omega)$. In particular $F_{k-1,k} \not\leq V(\omega)$. Next let $\mu \in \mathcal{W}_n$ correspond to the permutation $(k-1, k) \in \text{Sym}(n)$, and set $V := V(\omega)^\mu$. Then $V \leq U_n$ is a P_λ -suitable pattern group, and so, by Proposition 2.5, there is $\alpha \in \mathcal{W}_n$ that is good with respect to λ such that $V = V(\alpha)$.

As $F_{k,k-1} \not\leq V(\omega)$, $F_{k-1,l} \not\leq V(\omega)$ and $F_{k,l} \leq V(\omega)$, we have $F_{k-1,k} \not\leq V(\alpha)$, $F_{k,l} \not\leq V(\alpha)$ and $F_{k-1,l} \leq V(\alpha)$. That forces $\alpha(k) < \alpha(k-1)$, $\alpha(l) < \alpha(k)$ and $\alpha(k-1) < \alpha(l)$, respectively. This contradiction proves part (1). A similar argument proves part (2). \square

Corollary 2.12. *If $\omega(k) < \omega(k+1)$, then ω acts on the sets $\{1, \dots, k\}$ and $\{k+1, \dots, n\}$.*

Proof. Let $r \in \{1, \dots, k\}$. Then $\omega(r) < \omega(k+1)$, by Lemma 2.11 (1). Hence $\omega(r) < \omega(t)$, for all $t = k+1, \dots, n$, by Lemma 2.11 (2). In particular $\omega(\{1, \dots, k\}) \subseteq \{1, \dots, k\}$, and the statement follows. \square

Proposition 2.13. *Let $\omega \in \mathcal{W}_n$ be good with respect to λ such that $V(\omega)$ is P_λ -suitable. Then ω is a λ -permutation.*

Proof. First we show that $\omega = D_n(K_1, \dots, K_u)$, where $K_i \in \{I_{n_i}, \overline{I_{n_i}}\}$, and $n_1 + \dots + n_u = n$. Clearly there is some $k \in \{0, 1, \dots, n\}$ such that $\omega = D_n(K_1, \dots, K_u, \omega')$, where $K_i \in \{I_{n_i}, \overline{I_{n_i}}\}$, $n_1 + \dots + n_u = k$ and $\omega' \in \mathcal{W}_{n-k}$. Let us choose k maximal with this property and suppose $k < n$. Observe that ω acts on the sets $\{1, \dots, k\}$ and $\{k+1, \dots, n\}$. Also the maximality of k implies that $\omega(k+1) \neq k+1$. Next let $l \geq k+1$ be maximal such that $\omega(k+1) > \omega(k+2) \dots > \omega(l)$. Since $l = n$ or $\omega(l) < \omega(l+1)$, it follows from Corollary 2.12 that ω acts on the set $\{k+1, \dots, l\}$ and it does so like the permutation matrix $\overline{I_{l-k}}$. But this contradicts the maximality of k , and thus $k = n$.

Now let $i \in \{1, \dots, r\}$ so that $\lambda_i \geq 2$. As ω is good with respect to λ we have $\omega(s_i + 1) < \dots < \omega(s_{i+1})$. Now Corollary 2.12 implies that ω acts on the sets $\{1, \dots, s_i + 1\}$, $\{s_i + 2, \dots, s_{i+1} - 1\}$, $\{s_{i+1}, \dots, n\}$. In particular ω is a λ -permutation. \square

Theorem 2.14. *The set $\mathcal{C} := \{V(\omega) : \omega \in \mathcal{W}_n \text{ is a } \lambda\text{-permutation}\}$ provides a full set of representatives for the G -conjugacy classes of P_λ -suitable groups without repetitions.*

Proof. In Proposition 2.10 we have established that $V(\omega)$, where $\omega \in \mathcal{W}_n$ is a λ -permutation, is P_λ -suitable. Furthermore any two different such groups lie in different conjugacy classes, by Corollary 2.9.

Now let V be P_λ -suitable. Then V is G -conjugate to some $V' \leq U_n$. Hence $V' = U_n \cap P_\lambda^g$, for some $g \in G$. By Lemma 2.1 it follows that V' is G -conjugate to $V(\omega)$, where $\omega \in \mathcal{W}_n$ is good with respect to λ . In particular ω is a λ -permutation, by Proposition 2.13. \square

3. The number of components of $k_{P_\lambda} \uparrow^{\text{GL}_n}$ and their vertices

Let $\lambda = (\lambda_1, \dots, \lambda_r)$ be a composition of n . In the previous section we found all P_λ -suitable groups up to G -conjugation. In fact we have shown that V is P_λ -suitable if and only if $V =_G V(\omega)$, where ω is a λ -permutation. Hence the set \mathcal{C} from Theorem 2.14 contains a vertex for each component of $k_{P_\lambda} \uparrow^{\text{GL}_n}$. In this section we want to determine how many components have each of the groups in \mathcal{C} as a vertex. So for the remainder of this paper let $\omega \in \mathcal{W}_n$ be a λ -permutation and set $V := V(\omega)$.

By a result of Burry and Carlson [2] the kG -module $k_{P_\lambda} \uparrow^G$ and the $kN_G(V)$ -module $\bigoplus_{g \in A} k_{P_\lambda^g \cap N_G(V)} \uparrow^{N_G(V)}$ have the same number of components with vertex V , where A is a set of representatives g for those $(P_\lambda, N_G(V))$ -double cosets of G with $V \leq P_\lambda^g$.

Next suppose that $V \leq P_\lambda^g$, for some $g \in G$. Since U_n is a Sylow- p -group of P_λ it follows that $V^{g^{-1}h} \leq U_n$, for some $h \in P_\lambda$. Now by Lemma 2.8 we have $g^{-1}h \in N_G(V)$, and thus $g \in P_\lambda \cdot N_G(V)$. Hence by the above paragraph we obtain that $k_{P_\lambda} \uparrow^G$ has the same number of components with vertex V as $k_{P_\lambda \cap N_G(V)} \uparrow^{N_G(V)}$.

Lemma 3.1. *We have $P_\lambda \cap N_G(V(\omega)) = B_n$.*

Proof. Clearly $B_n \subseteq P_\lambda \cap N_G(V)$. Next let $X \in P_\lambda \cap N_G(V)$, such that $X \notin B_n$. Hence $X_{l,k} \neq 0$, for some $k < l$. Since $X \in N_G(V)$, Lemma 2.8 implies that there is some $j \in \{1, \dots, u\}$ such that $K_j = \overline{I_{n_j}}$ and $n_1 + \dots + n_{j-1} < k < l \leq n_1 + \dots + n_j$. In particular $\omega(k) > \omega(l)$.

On the other hand as $X \in P_\lambda$, then $s_i + 1 \leq k < l \leq s_{i+1}$, for some $i \in \{1, \dots, r\}$. But then $\omega(k) < \omega(l)$, as ω is good with respect to λ . This contradiction completes the proof. \square

Lemma 3.2. *Let $\omega \in \mathcal{W}_n$ be a λ -permutation. Then there is exactly one component in $k_{P_\lambda} \uparrow^{\text{GL}_n}$ with vertex $V(\omega)$.*

Proof. Let $V = V(\omega)$. By the introduction of this section and Lemma 3.1 we know that the number of components of $k_{P_\lambda} \uparrow^{\text{GL}_n}$ with vertex V coincides with the number of components of $k_{B_n} \uparrow^{N_G(V)}$ with vertex V . This number in turn equals the number of projective components of $k_{B_n/V} \uparrow^{N_G(V)/V}$. We have

$$B_n/V \cong \{D_n(A_1, \dots, A_u) : \text{where } A_i \in D_{n_i}, \text{ if } K_i = I_{n_i}, \\ \text{and } A_i \in B_{n_i}, \text{ if } K_i = \overline{I_{n_i}}\}$$

and

$$N_G(V)/V \cong \{D_n(A_1, \dots, A_u) : \text{where } A_i \in D_{n_i}, \text{ if } K_i = I_{n_i}, \\ \text{and } A_i \in \text{GL}_{n_i}, \text{ if } K_i = \overline{I_{n_i}}\},$$

where D_{n_i} is the group of diagonal matrices in GL_{n_i} . Hence

$$k_{B_n/V} \uparrow^{N_G(V)/V} \cong \bigotimes_{i: K_i = \overline{I_{n_i}}} k_{B_{n_i}} \uparrow^{\text{GL}_{n_i}}.$$

But every $k_{B_j} \uparrow^{\text{GL}_j}$ contains exactly one projective component, known as the Steinberg-module, (see [4] for more details). Now the statement follows by [6, Proposition 1.2]. \square

Finally we can state our main result.

Theorem 3.3. *The number of components of $k_{P_\lambda} \uparrow^{\text{GL}_n}$ coincides with the number of λ -permutations in \mathcal{W}_n , and $\{V(\omega) : \omega \in \mathcal{W}_n \text{ is a } \lambda\text{-permutation}\}$ gives a full set of the different vertices of the components of $k_{P_\lambda} \uparrow^{\text{GL}_n}$.*

Proof. We have seen that every component of $k_{P_\lambda} \uparrow^{\text{GL}_n}$ has a vertex of the form $V(\omega)$, where $\omega \in \mathcal{W}_n$ is a λ -permutation. By Corollary 2.9 we know that two such groups are not G -conjugate. The result of Lemma 3.2 completes the proof. \square

We conclude our paper with a specific example. Let $\lambda = (3, 1, 2)$ be a composition of $n = 6$. Then $s_1 = 0$, $s_2 = 3$, $s_3 = 4$ and $s_4 = 6$. Next observe that a λ -permutation ω acts on the sets $\{1\}$, $\{2\}$, $\{3, 4, 5\}$ and $\{6\}$. Hence there are exactly four λ -permutations and they are $\omega_1 = I_6$, $\omega_2 = D_6(I_2, \overline{I_3}, I_1)$, $\omega_3 = D_6(I_2, \overline{I_2}, I_2)$ and $\omega_4 = D_6(I_3, \overline{I_2}, I_1)$. In particular $k_{P_\lambda} \uparrow^{\text{GL}_6}$ has exactly four components with the respective vertices

$$\begin{aligned} V(\omega_1) &= U_6 \\ V(\omega_2) &= \{X \in U_6 : X_{3,4} = X_{3,5} = X_{4,5} = 0\}, \\ V(\omega_3) &= \{X \in U_6 : X_{3,4} = 0\}, \\ V(\omega_4) &= \{X \in U_6 : X_{4,5} = 0\}. \end{aligned}$$

References

- [1] A. Borel: *Linear Algebraic Groups*, Springer-Verlag, New York, 1991.
- [2] D.W. Burry and J.F. Carlson: *Restriction of modules to local subgroups*, Proc. Amer. Math. Soc. **84** (1982), 181–184.
- [3] M. Cabanes: *Brauer morphism between modular Hecke algebras*, J. Algebra **115** (1988), 1–31.
- [4] J.E. Humphreys: *Modular Representations of Finite Groups of Lie Type*, London Mathematical Society Lecture Note Series **326**, Cambridge University Press, Cambridge, 2006.
- [5] I.M. Isaacs: *Counting characters of upper triangular groups*, J. Algebra **315** (2007), 698–719.
- [6] B. Külshammer: *Some indecomposable modules and their vertices*, J. Pure Appl. Algebra **86** (1993), 65–73.
- [7] L. Pforte: *A technique for the search for vertices of the components of a kG -permutation module*, Comm Algebra **43** (2015), 659–673.
- [8] P. Sin: *The Green ring and modular representations of finite groups of Lie type*, J. Algebra **123** (1989), 185–192.
- [9] N. Tinberg: *Some indecomposable modules for finite groups with split BN-pairs*, J. Algebra **61** (1979), 508–526.

Department of Mathematics
National University of Ireland, Maynooth
Co. Kildare
Ireland
e-mail: larspforte@yahoo.ie